

1     **Experimental transmission of the chronic wasting disease agent to**  
2                     **swine after oral or intracranial inoculation**

3     Running Title: The chronic wasting disease agent transmits to swine

4  
5  
6     S. Jo Moore<sup>1,2</sup>, M. Heather West Greenlee<sup>3</sup>, Naveen Kondru<sup>3</sup>, Sireesha Manne<sup>3</sup>, Jodi D.  
7     Smith<sup>1,#</sup>, Robert A. Kunkle<sup>1</sup>, Anumantha Kanthasamy<sup>3</sup>, Justin J. Greenlee<sup>1\*</sup>

8  
9     <sup>1</sup> Virus and Prion Research Unit, National Animal Disease Center, USDA, Agricultural  
10    Research Service, Ames, Iowa, United States of America

11  
12    <sup>2</sup> Oak Ridge Institute for Science and Education, Oak Ridge, Tennessee, United States of  
13    America

14  
15    <sup>3</sup> Department of Biomedical Sciences, Iowa State University College of Veterinary  
16    Medicine, Ames, Iowa, United States of America

17  
18    <sup>#</sup> Current Address: Department of Veterinary Pathology, Iowa State University College of  
19    Veterinary Medicine, Ames, Iowa, United States of America

20  
21    \* Corresponding author

22    Email: [justin.greenlee@ars.usda.gov](mailto:justin.greenlee@ars.usda.gov)

23

## Abstract

Chronic wasting disease (CWD) is a naturally occurring, fatal neurodegenerative disease of cervids. The potential for swine to serve as a host for the agent of chronic wasting disease is unknown. The purpose of this study was to investigate the susceptibility of swine to the CWD agent following experimental oral or intracranial inoculation. Crossbred piglets were assigned to one of three groups: intracranially inoculated (n=20), orally inoculated (n=19), or non-inoculated (n=9). At approximately the age at which commercial pigs reach market weight, half of the pigs in each group were culled ('market weight' groups). The remaining pigs ('aged' groups) were allowed to incubate for up to 73 months post inoculation (MPI). Tissues collected at necropsy were examined for disease-associated prion protein (PrP<sup>Sc</sup>) by western blotting (WB), antigen-capture immunoassay (EIA), immunohistochemistry (IHC) and *in vitro* real-time quaking induced conversion (RT-QuIC). Brain samples from selected pigs were also bioassayed in mice expressing porcine prion protein. Four intracranially inoculated aged pigs and one orally inoculated aged pig were positive by EIA, IHC and/or WB. Using RT-QuIC, PrP<sup>Sc</sup> was detected in lymphoid and/or brain tissue from one or more pigs in each inoculated group. Bioassay was positive in 4 out of 5 pigs assayed. This study demonstrates that pigs can support low-level amplification of CWD prions, although the species barrier to CWD infection is relatively high. However, detection of infectivity in orally inoculated pigs using mouse bioassay raises the possibility that naturally exposed pigs could act as a reservoir of CWD infectivity.

## 46 **Importance**

47       We challenged domestic swine with the chronic wasting disease agent by  
48 inoculation directly into the brain (intracranially) or by oral gavage (orally). Disease-  
49 associated prion protein (PrP<sup>Sc</sup>) was detected in brain and lymphoid tissues from  
50 intracranially and orally inoculated pigs as early as 8 months of age (6 months post-  
51 inoculation). Only one pig developed clinical neurologic signs suggestive of prion  
52 disease. The amount of PrP<sup>Sc</sup> in the brains and lymphoid tissues of positive pigs was  
53 small, especially in orally inoculated pigs. Regardless, positive results in orally  
54 inoculated pigs suggest that it may be possible for swine to serve as a reservoir for prion  
55 disease under natural conditions.

56

## 57 **Introduction**

58       The transmissible spongiform encephalopathies (TSEs) or prion diseases are fatal  
59 neurodegenerative diseases. Naturally occurring TSEs include chronic wasting disease  
60 (CWD) in cervids, scrapie in sheep, bovine spongiform encephalopathy (BSE) in cattle,  
61 and sporadic and familial prion diseases in humans.

62       The potential for swine to serve as a host for the agent of CWD is unknown. A  
63 naturally occurring TSE has not been reported in swine (1, 2). Intracranial challenge of  
64 swine with kuru, a human prion disease, was not successful (3), although, at the time of  
65 those studies, molecular tests for disease-associated prion protein (PrP<sup>Sc</sup>) were not  
66 available. Pigs have been shown to be susceptible to bovine BSE following parenteral  
67 inoculation (simultaneous intraperitoneal, intravenous and intracranial routes), ovine BSE  
68 following intracranial inoculation (4), and ovine scrapie following intracranial

69 inoculation (5), but not to bovine BSE after oral challenge with large amounts of infected  
70 brain material (6-8).

71         The CWD agent has a wide host range amongst cervids and can be experimental  
72 transmitted to several other species. Naturally occurring CWD has been reported in  
73 cervids including mule deer (*Odocoileus hemionus*) (9-11), Rocky mountain elk (*Cervus*  
74 *elaphus nelson*) (11, 12), white-tailed deer (*Odocoileus virginianus*) (10, 11), moose  
75 (*Alces alces shirasi*) (13, 14), and reindeer (*Rangifer tarandus tarandus*) (15). In  
76 addition, Eurasian red deer (*Cervus elaphus*) (16), Eurasian fallow deer (*Dama dama*)  
77 (17), Asian muntjac (*Muntiacus reevesi*) (18), and reindeer (19, 20) have been shown to  
78 be susceptible to CWD following experimental inoculation. CWD has been  
79 experimentally transmitted to non-cervid species including sheep (21), cattle (22-25),  
80 domestic cats (26, 27), ferrets (28, 29), non-human primates (30-32), and laboratory  
81 rodents (reviewed in (33)).

82         Pigs could be exposed to CWD infectivity via two main routes: 1) Exposure of  
83 farmed or pet swine (*Sus scrofa domesticus*) to contaminated feed and 2) exposure of  
84 feral swine (*Sus scrofa*) to CWD infected carcasses or contaminated environments. In the  
85 US, feeding of ruminant by-products to ruminants is prohibited, but feeding of ruminant  
86 materials to swine, mink, and poultry still occurs. Therefore, it is possible that, if a CWD-  
87 affected cervid carcass entered the food chain through a commercial slaughter house,  
88 domesticated farmed and pet swine could be exposed to CWD infectivity in  
89 commercially prepared rations. As of 2015, feral pigs have been reported in 39 US states  
90 (34) and in 12 of these states CWD has been detected in free-ranging cervid populations  
91 (35). Environmental contamination with CWD infectivity in excreta or decomposing

92 carcasses contributes to horizontal transmission of CWD in mule deer (10). Prion  
93 infectivity has been shown to persist on the surface of contaminated plant leaves and  
94 roots (36) and in soil (37-39). Therefore, feral pigs could be exposed to infectivity  
95 through scavenging of CWD-affected carcasses, consumption of contaminated  
96 vegetation, and while rooting around in the soil during foraging. In this study we  
97 demonstrate that swine are susceptible to the CWD agent following oral or intracranial  
98 experimental inoculation and accumulate PrP<sup>Sc</sup> in both brain and lymphoid tissues.  
99 Detection of PrP<sup>Sc</sup> in brain and lymphoid tissues from orally inoculated pigs at 6 months  
100 after inoculation raises the possibility that naturally exposed pigs could potentially be a  
101 reservoir for CWD prions.

102

## 103 **Results**

### 104 **Clinical presentation**

105 All pigs culled at 6 MPI (8 months of age; n = 8 intracranially (IC) inoculated, n =  
106 9 orally inoculated) were clinically normal with the exception of one pig (#35) that was  
107 noted to be limping on its left front and rear legs. Four IC inoculated pigs and 1 orally  
108 inoculated pig) developed intercurrent lameness from approximately 30 MPI usually  
109 beginning with the feet and legs and progressing to difficulty rising. At approximately 41  
110 MPI 4 clinically normal pigs (n = 1 IC inoculated, n = 4 orally inoculated) were culled to  
111 reduce animal density in the containment space. Neurological signs were observed in one  
112 pig (animal #27, incubation period = 64 MPI) and included difficulty rising, and muscle  
113 fasciculations and tremors after rising. Pig #27 also had skin abrasions and/or ulceration

114 over pressure points and polyarthritis. All other pigs were found dead or culled due to  
115 intercurrent disease, most commonly lameness that was not responsive to treatment.

116

#### 117 **Detection of PrP<sup>Sc</sup>**

118 To determine if pigs inoculated with the agent of CWD accumulate misfolded  
119 prion protein in the central nervous system we assayed brainstem using western blot  
120 (WB), enzyme immunoassay (EIA), immunohistochemistry (IHC) and *in vitro* real-time  
121 quaking induced conversion (RT-QuIC). Results of screening of brainstem material from  
122 all pigs by WB and EIA, and results for additional testing of animals that were PrP<sup>Sc</sup>  
123 positive by either screening test can be found in Table 1.

124

#### 125 *Western blotting*

126 Using WB, PrP<sup>Sc</sup> was detected in brain tissue from two intracranially inoculated  
127 pigs (#27, #28) necropsied at 64 and 73 MPI, respectively (Table 1).

128 The migration pattern of samples from pigs inoculated IC with the CWD agent were  
129 different from either the sample from a pig inoculated with classical bovine spongiform  
130 encephalopathy (BSE) or the original CWD inoculum (Figure 1). While the  
131 monoglycosylated (middle) band was most prominent in the sample from the pig  
132 inoculated with BSE, the diglycosylated (top) band was most prominent in the sample  
133 from the pig inoculated with the CWD agent and the original CWD inoculum.

134

#### 135 *Enzyme immunoassay*

136 Using EIA, misfolded protein was detected in brain tissue from 1/10 IC  
137 inoculated market weight pigs, 5/10 IC inoculated aged pigs (42-73 MPI), 0/9 orally  
138 inoculated market weight pigs, and 1/10 orally inoculated aged pig (Table 1).

139

#### 140 *Real-time quaking induced conversion*

141 Using RT-QuIC, PrP<sup>Sc</sup> was detected in brainstem material from 3/6 IC inoculated  
142 market weight pigs, 7/7 IC inoculated aged pigs, 2/6 orally inoculated market weight  
143 pigs, and 5/6 orally inoculated aged pigs (Table 1, Figure 2). For each positive sample we  
144 quantified the seeding activity based on amyloid formation rate (AFR), which is the  
145 reciprocal of the time (h) that it takes for a reaction to reach the threshold (Ct), defined as  
146 the mean baseline fluorescence plus 5 standard deviations. For IC inoculated pigs (n =  
147 10), the mean AFR for each animal ranged from 0.025-0.210. For orally inoculated pigs  
148 (n = 7) the range of mean AFRs was 0.010-0.029 (Table 1, Figure 2). Average RT-QuIC  
149 data, generated by calculating the mean of all replicates from all animals in each  
150 challenge group, can be found in Figure 3.

151

#### 152 **Differential proteinase K (PK) sensitivity of brainstem samples**

153 To investigate possible biochemical properties of PrP<sup>Sc</sup> that may have contributed  
154 to the variation in aggregation kinetics observed on the RT-QuIC assay, the EIA optical  
155 density was measured on matched samples with and without treatment with PK. The  
156 difference in optical density between non-PK treated and PK treated samples allows us to  
157 estimate the relative PK resistance of the PrP<sup>Sc</sup> present in the brains of infected pigs (40).

158 PrP<sup>Sc</sup> in EIA positive brain tissue from one IC inoculated market weight pig  
159 (#15), one orally inoculated aged pig (#45), and one IC inoculated aged pig (#24) was PK  
160 sensitive. PrP<sup>Sc</sup> from the remaining 4 pigs with samples positive by EIA, all from the IC  
161 inoculated aged pig group, was PK resistant (Table 1). Proteinase-K titration was  
162 performed on all EIA positive samples and results were consistent across PK  
163 concentrations of 0.4-50 µg/mL.

164 Six brain samples were EIA positive and RT-QuIC positive. Of these, the 4  
165 samples that were PK resistant had higher AFRs (range = 0.17-0.21), while the 2 samples  
166 that were PK sensitive had lower AFRs (0.01 and 0.03) (Figure 2).

167

#### 168 **Detection of PrP<sup>Sc</sup> in lymphoid tissues**

169 To determine if pigs inoculated with the CWD agent accumulate misfolded prion  
170 protein in lymphoid tissues, EIA and RT-QuIC were applied to samples of the  
171 retropharyngeal lymph node (RPLN), palatine tonsil, and mesenteric lymph node (MLN).  
172 Full results for individual pigs can be found in Table 2.

173 All lymphoid tissues tested were PrP<sup>Sc</sup> negative by EIA with the exception of pig  
174 #37 (orally inoculated market weight pig) that had a positive MLN. Using the RT-QuIC  
175 assay, PrP<sup>Sc</sup> was detected in lymphoid tissues of the head (RPLN, palatine tonsil) in 3/6  
176 IC inoculated market weight pigs, 5/7 IC inoculated aged pigs, 4/6 orally inoculated  
177 market weight pigs, and 2/6 orally inoculated aged pigs. The MLN was positive in 5/6  
178 orally inoculated market weight pigs, 3/4 orally inoculated aged pigs (samples were not  
179 available for 2 pigs), 4/6 IC inoculated market weight pigs, 2/4 IC inoculated aged pigs.



180 Overall, the MLN was positive in 14/19 (74%) of samples examined, the RPLN in 8/18  
181 (44%), and the tonsil in 10/25 (40%).

182

### 183 **Histopathology and Immunohistochemistry**

184 To determine if pigs inoculated with the CWD agent develop spongiform lesions  
185 or accumulate misfolded prion protein in the brain, coronal brain sections were examined  
186 by light microscopy after H&E staining and immunohistochemistry.

187 Occasional neuropil vacuolation and white matter vacuolation were present in different  
188 brain sections of control and inoculated pigs. Small to medium-sized grey matter  
189 vacuoles were seen in the colliculus of at least one pig from each treatment group,  
190 including control pigs (Figure 4A, pig #7 and Figure 4B, #25). Vacuolation and PrP<sup>Sc</sup>  
191 deposition in the colliculus was present in two pigs (#25, #26) from the IC inoculated  
192 aged pig group (Figure 4C, #25). Intraneuronal vacuolation was observed in large  
193 neurons of the dorsal motor nucleus of the vagus nerve (DMNV) in the medulla at the  
194 level of the obex (Figure 4E, #38). This type of vacuolation was present in pigs from all  
195 market weight treatment groups including non-inoculated control pigs, and in aged  
196 control pigs. PrP<sup>Sc</sup> deposition in association with DMNV vacuolation was not observed in  
197 any pigs.

198

199 Positive PrP<sup>Sc</sup> immunoreactivity was observed in samples from 4 pigs. In the brain, PrP<sup>Sc</sup>  
200 immunoreactivity appeared as the intraneuronal type (coarse granular deposits of PrP<sup>Sc</sup> in  
201 the neuronal perikarya surrounding the nucleus) in large neurons of the rostral medulla

202 reticular formation (#26), midbrain colliculus (#25, #26), midline thalamic nuclei and  
203 hypothalamus (#45, #28), or septal nuclei (#28) (Figure 4C).

204 PrP<sup>Sc</sup> immunoreactivity was also seen in the retina of one pig (#26), granular to  
205 punctate immunoreactivity in the inner and out plexiform layers with occasional intragial  
206 deposits (Figure 4F, #26). Disease-specific PrP<sup>Sc</sup> immunoreactivity was not seen in any  
207 other tissues although non-specific immunolabeling was common (Figure 4D, brainstem;  
208 Figure 4G, retina).

209

#### 210 **Mouse bioassay**

211 To determine if pigs inoculated with the CWD agent accumulate infectious  
212 material, brainstem material from selected pigs was bioassayed in Tg002 mice that  
213 express porcine prion protein at normal levels (5).

214 Pigs from the IC inoculated market weight (#18) and IC inoculated aged (#27,  
215 #28) groups, and the orally inoculated aged group (#48) produced positive bioassay  
216 results (Table 3). In mice inoculated with brain material from pig #18 (IC inoculated  
217 market weight pig) the average incubation period was 244 days post-inoculation (dpi)  
218 (2/28 mice). In mice inoculated with brain material from pig #27 (IC inoculated market  
219 aged pig group) the average incubation period was 167 dpi (3/29 mice, range 140-220  
220 dpi). Two out of 27 mice were positive in the group inoculated with brain material from  
221 pig #28; one mouse was found dead at 314 dpi and the other was euthanized at the end of  
222 the study at 701 dpi. The highest attack rate resulted from the orally inoculated aged pig  
223 (#48), with 14/28 mice positive and an average incubation period of 263 dpi (range 111-  
224 621 dpi).

225 All pigs that produced a positive bioassay result also had a positive RT-QuIC  
226 result. In addition, pig #27 and #28 were positive by WB (both pigs), EIA (both pigs),  
227 and IHC (#28 only). Bioassay of brain tissue from pig #32 in the orally inoculated market  
228 weight group was unsuccessful (0/28 mice, study ended at 702 dpi) (Table 3), although  
229 PrP<sup>Sc</sup> was detected in the brain of this pig using RT-QuIC (Table 1).

## 231 Discussion

232 We demonstrated that PrP<sup>Sc</sup> can be detected in brain and lymphoid tissues from as  
233 early as 6 months post-inoculation in pigs inoculated orally or intracranially with the  
234 CWD agent. We show that pigs inoculated with CWD rarely develop neurologic signs  
235 suggestive of prion disease although PrP<sup>Sc</sup> can be detected in brain samples. Furthermore,  
236 neuropathological changes are often equivocal and the amount of PrP<sup>Sc</sup> present is  
237 generally low, so sensitive methods such as real-time quaking induced conversion (RT-  
238 QuIC) and bioassay were used for PrP<sup>Sc</sup> detection.

239 Prion infection was subclinical in most pigs in this study; PrP<sup>Sc</sup> was detected in  
240 brain from 18 pigs but neurologic signs suggestive of prion disease were observed in only  
241 one pig. This pig developed clinical signs of difficulty in rising and signs of tremor. Both  
242 these clinical signs have been reported previously in pigs challenged with bovine BSE (6)  
243 or sheep-passaged BSE (4). A number of pigs developed persistent recumbency with  
244 difficulty in rising, but these clinical signs were attributed to musculoskeletal lameness  
245 rather than neurological disease.

246 Similar to pigs with BSE (8), PrP<sup>Sc</sup> accumulation was sparse and did not  
247 necessarily correlate with the degree of spongiform change. In addition to having a

248 restricted distribution, the range of morphological types of PrP<sup>Sc</sup> was limited to just the  
249 intraneuronal type. Prominent intraneuronal immunolabeling is also a feature of scrapie  
250 in pigs (5). In contrast, a wider variety of PrP<sup>Sc</sup> deposit types have been described in pigs  
251 challenged with bovine (6, 8) or sheep-passaged BSE (4).

252 Mild spongiform change was observed in the brain of both inoculated and non-  
253 inoculated pigs, suggesting that the presence of spongiform change in the brain should  
254 not be used as a sole diagnostic test for CWD in pigs. Similar to results reported by  
255 others, microscopic changes in negative control and inoculated pigs were limited to  
256 occasional scattered vacuoles in the neuropil or white matter throughout the brain (1),  
257 neuropil vacuolation of the superficial layers of the rostral colliculus (1, 8), and  
258 occasional neuronal vacuolation in the dorsal motor nucleus of the vagus nerve (1, 2, 8).  
259 Since the above microscopic changes can be observed in both non-inoculated control and  
260 inoculated pigs, when present in inoculated pigs they are considered equivocal, i.e. not  
261 related to prion disease. Co-localization of neuropil vacuolation and intraneuronal PrP<sup>Sc</sup>  
262 deposits were present in the rostral colliculus of 2 pigs in our study but vacuolation did  
263 not extend to deeper layers of the rostral colliculi or to other areas of the brain (8), so was  
264 considered equivocal.

265 Limited microscopic and immunohistopathological changes observed in the brains  
266 of pigs with CWD compared to pigs inoculated with bovine or ovine-adapted BSE  
267 suggests that the species barrier for CWD to pigs is higher than for BSE to pigs. Despite  
268 this, pigs are able to accumulate misfolded prion protein and CWD infectivity.

269 Using standard diagnostic tests (western blot, enzyme immunoassay, or  
270 immunohistochemistry), PrP<sup>Sc</sup> was detected in brain or lymphoid tissues from 8 pigs

271 from this study. The number of positive animals and tissues, in particular lymphoid  
272 tissues, was much higher when the RT-QuIC assay was used. Using RT-QuIC, PrP<sup>Sc</sup> was  
273 detected in brain and lymphoid tissues that were PrP<sup>Sc</sup> negative by all other tests. This is  
274 not surprising considering that RT-QuIC is reported to be at least as sensitive as bioassay  
275 (41) and 10,000 fold more sensitive than enzyme immunoassay and western blot assays  
276 for the detection of scrapie seeding activity in goat brain samples (42). With the  
277 exception of immunohistochemistry, diagnostic tests were performed on brainstem  
278 samples since this brain region is the preferred site for statutory diagnostic testing.  
279 Testing of additional brain regions may have revealed PrP<sup>Sc</sup> accumulation elsewhere in  
280 the brain, as was observed on immunohistochemistry.

281         The RT-QuIC assay allows quantification of the seeding activity of prions in the  
282 samples based upon amyloid formation rate (AFR) values. The AFR is calculated as the  
283 reciprocal of the time taken for a reaction to reach the threshold (i.e. 1/(time to threshold  
284 in hours)). A higher AFR reflects a shorter time taken to threshold, which can also be  
285 termed a shorter 'lag phase'. Lag phases have previously been shown to be inversely  
286 correlated with seed concentration in RT-QuIC reactions (41, 43, 44). Since the AFRs of  
287 samples from IC inoculated aged pigs tended to be higher than those from orally  
288 inoculated aged pigs, it follows that the relative amount of PrP<sup>Sc</sup> in the brain is higher in  
289 IC inoculated pigs. This seems logical considering that PrP<sup>Sc</sup> in the inoculum was  
290 delivered directly into the brain in IC inoculated pigs, but delivered to peripheral tissues  
291 (oral cavity and gastrointestinal tract) in orally inoculated pigs.

292         We observed that the AFR of samples from positive animals that were determined  
293 to be PK sensitive was approximately one order of magnitude lower than the AFR of

294 samples that were PK resistant. Although the interpretation of these observations is  
295 limited by the small sample size and the fact that samples were not normalized for total  
296 protein content, it appears that there may be a relationship between AFR and PK  
297 sensitivity.

298         One hypothesis is that larger seed particles present more seeding surfaces than  
299 smaller particles and thus support faster RT-QuIC kinetics (45). In scrapie infected  
300 hamsters, PK sensitive PrP<sup>Sc</sup> molecules from low molecular weight aggregates are made  
301 up of fewer PrP units (i.e. are smaller) than PK resistant PrP<sup>Sc</sup> aggregates (46, 47).  
302 Combining these observations with our own results, we hypothesize that the smaller  
303 average seed particle size of PK sensitive PrP<sup>Sc</sup> may result in slower RT-QuIC kinetics  
304 and leads to lower AFRs and longer lag times. However, as stated above, this hypothesis  
305 is based on a small number of samples.

306  
307         The detection of PrP<sup>Sc</sup> in lymphoid tissues from the head and gut of CWD-  
308 infected pigs raises the possibility that pigs may be able to shed prions in excreta as has  
309 been shown for saliva (48-51) and feces (52-54) from CWD-affected cervids.  
310 Unfortunately, saliva and feces were not collected in the current study.

311         PrP<sup>Sc</sup> was detected in brain and lymphoid tissues from orally inoculated pigs  
312 killed at approximately market weight. These results suggest that, if they were to be  
313 exposed to sufficient amounts of CWD infectivity, pigs in commercial swine production  
314 systems have the potential to accumulate CWD prions by the time they reach market  
315 weight.

316           In the case of feral pigs, exposure to the agent of CWD through scavenging of  
317 CWD-affected cervid carcasses or through consumption of prion contaminated plants or  
318 soil could allow feral pigs to serve as reservoirs of CWD infectivity. The range and  
319 numbers of feral pigs is predicted to continue to increase due to the ability of pigs to  
320 adapt to many climates, reproduce year-round, and survive on a varied diet (55). The  
321 range of CWD-affected cervids also continues to spread, increasing the likelihood of  
322 overlap of ranges of feral pigs and CWD-affected environments.

323

324           We demonstrate here that PrP<sup>Sc</sup> accumulates in lymphoid tissues from pigs  
325 inoculated intracranially or orally with the CWD agent, and can be detected as early as 6  
326 months after inoculation. Clinical disease suggestive of prion disease developed only in a  
327 single pig after a long (64 months) incubation period. This raises the possibility that  
328 CWD-infected pigs could shed prions into their environment long before they develop  
329 clinical disease. However, the low amounts of PrP<sup>Sc</sup> detected in the study pigs combined  
330 with the low attack rates in Tg002 mice suggest that there is a relatively strong species  
331 barrier to CWD prions in pigs.

## 332 **Materials and methods**

### 333 **Ethics statement**

334 All animal experiments were reviewed and approved by the National Animal  
335 Disease Center's (NADC) Institutional Animal Care and Use Committee (protocol  
336 numbers 3510 (swine) and 2422 (mice)) and were carried out in strict accordance with  
337 the Guide for the Care and Use of Laboratory Animals (Institute of Laboratory Animal  
338 Resources, National Academy of Sciences, Washington, DC) and the Guide for the Care  
339 and Use of Agricultural Animals in Research and Teaching (Federation of Animal  
340 Science Societies, Champaign, IL). Pigs were observed daily for clinical signs of disease  
341 and euthanized and necropsied at approximately 6 months post-inoculation, or when  
342 unequivocal signs of prion disease such as behavior changes, decreased feed intake, loss  
343 of body condition, ataxia, prolonged recumbency, or inability to rise were confirmed by a  
344 veterinarian, or when euthanasia was necessary due to intercurrent illness or injury that  
345 could not be remediated by veterinary care. Euthanasia was performed by intravenous  
346 injection of sodium pentobarbital according to the manufacturer's instructions.

347

### 348 **Inoculum preparation**

349 The pooled CWD inoculum was prepared from 3 brains from white-tailed deer  
350 that were inoculated intracranially with brain material from CWD-affected elk, white-  
351 tailed deer or mule deer (NADC Institutional Animal Care and Use Committee, protocol  
352 number 3347) (56). All donor deer were homozygous for glycine (G/G) at *PRNP* codon  
353 96, and homozygous for serine (S/S) at codon 138. The brain tissue was ground in a



354 mechanical grinder and mixed with phosphate-buffered saline to produce a 10% weight  
355 per volume (w/v) homogenate.

356

### 357 **Animal procedures**

358 Crossbred piglets were inoculated at 8 weeks of age. Intracranially inoculated pigs  
359 (n = 20) received a single intracranial inoculation of 0.75 mL of 10% w/v CWD brain  
360 homogenate as described previously (57). Orally inoculated pigs (n = 19) received 15 mL  
361 of 10% w/v CWD brain homogenate by syringe with a soft feeding tube on four  
362 consecutive days (total dose 45 mL). Intracranially and orally inoculated pigs were  
363 housed in separate pens. At 2 weeks post-inoculation non-inoculated control pigs were  
364 introduced into the pens with the inoculated pigs.

365 At 6-7 months of age, approximately the time at which commercial pigs reach  
366 market weight, half of the pigs in each group were culled ('market weight' groups): n = 8  
367 intracranially inoculated pigs, n = 9 orally inoculated pigs, and n = 2 control pigs. The  
368 remaining pigs ('aged' groups) were allowed to incubate for up to 73 months post  
369 inoculation (MPI) when the study ended. Swine were observed daily for the development  
370 of clinical signs.

371

### 372 **Mouse bioassay**

373 Infectivity in brain tissue from selected pigs was assayed via intracranial  
374 inoculation of *Tg002* mice that express porcine prion protein (GenBank porcine sequence  
375 accession no. GU595061) at approximately 1X the expression level of prion protein in  
376 FVB mice (5). Samples of brainstem at the level of the obex were prepared as 10% w/v

377 homogenates in PBS. Mice were inoculated intracranially with 20  $\mu$ L of 10% w/v brain  
378 homogenate as described previously (58). Mice were monitored daily and euthanized  
379 when they displayed unequivocal neurological signs (difficulty moving, poor  
380 coordination, unable to move, anorexia) or at the time of study termination  
381 (approximately 700 days post-inoculation). Brain samples from mice were prepared as  
382 10% w/v brain homogenates in phosphate buffered saline as described previously (59).  
383 PrP<sup>Sc</sup> was detected using enzyme immunoassay as described below.

384

#### 385 **Sample collection**

386 A full necropsy was performed on all pigs including collection of two sets of  
387 tissue samples. To minimize potential cross-contamination one pathologist collected  
388 tissues from the head and a second pathologist collected tissues from the rest of the body.  
389 Single use instruments were not used. One set of tissues included representative sections  
390 of liver, kidney, spleen, skin, striated muscles (heart, tongue, diaphragm, masseter,  
391 triceps, biceps femoris, psoas major), lymphoid tissues of the head (pharyngeal tonsil,  
392 palatine tonsil, medial retropharyngeal lymph node), other lymph nodes (mesenteric,  
393 hepatic, renal, popliteal, prescapular), nasal turbinates, lung, esophagus, small intestine,  
394 cecum, colon, rectal mucosa, stomach, adrenal gland, pituitary gland, reproductive  
395 tissues, peripheral nervous system (trigeminal ganglion, optic nerve, sciatic nerve, vagus  
396 nerve), brain (hemisections of cerebral cortex, hippocampus, cerebellum, superior  
397 colliculus and brainstem including obex) and eye (retina). Formalin-fixed tissues were  
398 fixed in 10% neutral buffered formalin, moved to 70% ethyl alcohol after 48 hours,

399 embedded in paraffin wax, sectioned, and stained with hematoxylin and eosin (HE) for  
400 light microscopy. The second set of tissues was frozen.

401

#### 402 **Selection of animals and tissues for PrP<sup>Sc</sup> detection**

403 Frozen brainstem from all pigs was screened for the presence of PrP<sup>Sc</sup> using  
404 antigen-capture enzyme immunoassay (EIA) and western blot (WB). Fixed tissues from  
405 pigs that were positive on WB and/or EIA were examined by immunohistochemistry  
406 (IHC). In addition, representative pigs from across the range survival times for each  
407 group were also examined by IHC. Brainstem material from the pig with the longest  
408 incubation period in each treatment group was bioassayed in *Tg002* mice. PrP<sup>Sc</sup> detection  
409 using quaking-induced conversion assay (QuIC) was applied to frozen brainstem and  
410 lymphoid tissue from all pigs that were positive by any other test (EIA, WB, IHC,  
411 bioassay) as well as additional animals so that 6-7 animals per group and across a range  
412 of survival times were tested.

413

#### 414 **Immunohistochemistry**

415 All paraffin-embedded tissues were immunostained by an automated  
416 immunohistochemical method for detection of PrP<sup>Sc</sup> as described previously (60) using  
417 the anti-PrP monoclonal antibody L42.

418

#### 419 **Antigen-capture enzyme immunoassay (EIA)**

420 Brain homogenates were homogenized in 1X PBS at a concentration of 20% w/v  
421 and assayed with a commercially available EIA kit (HerdChek BSE-Scrapie Ag Test Kit,

422 EIA, IDEXX Laboratories, Westbrook, ME) as previously described (61). Assays were  
423 performed in accordance with the manufacturer's instructions. The EIA kit instructions  
424 indicated 3 protocols (standard, short, and ultrashort). The short protocol was used for  
425 testing of tissue samples in the present study. Each tissue sample homogenate was  
426 assayed in a single well along with negative and positive controls supplies with the kit.  
427 Two conjugate concentrate products were included with the kit: a conjugate concentrate  
428 intended for use with brain samples obtained from small ruminants (SRB-CC) and a  
429 conjugate concentrate intended for use with brain samples obtained from cattle or lymph  
430 node or spleen samples obtained from small ruminants (CC). In this study (SRB-CC)  
431 conjugate was used for testing the samples obtained from mice expressing pig prion  
432 protein. Absorbance was measured (SPECTRAmax 190, Molecular Devices, Sunnyvale,  
433 CA) at 450 nm with a reference wavelength of 620 nm. Cutoff values were established  
434 for each run according to kit instructions whereby 0.180 was added to the mean negative  
435 control value. Samples were interpreted as positive if their absorbance value at 450 nm  
436 minus the reference value at 620 nm was above the established cutoff value.

437

#### 438 **EIA-based proteinase-K sensitivity testing**

439 Sensitivity to proteinase-K (PK) was determined using the EIA protocol described  
440 above but with the addition of a pre-testing PK-treatment step (40). Briefly, for each  
441 animal two 100  $\mu$ L aliquots of 20% w/v brain homogenate were prepared: 5  $\mu$ L of 1  
442 mg/mL PK (USB Corporation, Cleveland, OH, USA) was added to one aliquot and 5  $\mu$ L  
443 of PBS was added to the second aliquot. Both aliquots were incubated for 1 hour at 37 °C  
444 with shaking at 1000 rpm followed by the addition of 1.0  $\mu$ L of 100mg/mL PK-inhibitor

445 (Pefabloc, Roche Diagnostics, Mannheim, Germany). The absorbance value for each  
446 sample was determined using EIA as described above. Samples for which the non-PK  
447 treated aliquot was EIA positive and the PK-treated aliquot was EIA negative were  
448 classified as PK sensitive. Samples for which the non-PK treated aliquot was EIA  
449 positive and the PK-treated aliquot was EIA positive were classified as PK-resistant.

450

#### 451 **Western blotting**

452 Samples for WB were collected from the brainstem at the level of the obex and  
453 the midbrain between the optic and oculomotor nerves dorsal to the pituitary and  
454 performed as previously reported (57). Tissues were homogenized and enriched as  
455 described previously (22) with the following modifications: After the pellets were  
456 resuspended in 100  $\mu$ L of water, samples were digested with proteinase K (PK) using a  
457 final enzyme concentration of 0.4 U/mL (8  $\mu$ g/mL) at 37 °C for 1 hr. The digestion was  
458 stopped by the addition of a serine protease inhibitor (Pefabloc SC, Roche Diagnostics  
459 GmbH, Mannheim, Germany) to a final concentration of 1 mg/mL. Western blots were  
460 developed using mouse anti-PrP monoclonal antibody L42, which targets to amino acids  
461 145-163 of the ovine prion protein sequence (62), at 1:500 dilution (0.1  $\mu$ g/mL).

462 Due to the sparse PrP<sup>Sc</sup> accumulation in the brains of inoculated pigs the blot in Figure 1  
463 is a composite. The Pig CWD sample was enriched and loaded at 100 mg/eq. The Pig  
464 BSE positive control tissue was provided by the APHA Biological Archive (Addlestone,  
465 UK).

466

#### 467 **Expression and purification of the recombinant PrP substrate**

468           The recombinant prion protein (rPrP) used in the RT-QuIC assay was expressed  
469 and purified using a standard protocol from previous reports (41, 63). Briefly, rPrP  
470 composed of Syrian hamster PrP residues 90-231 in the pET vector was transformed into  
471 *Escherichia coli* Rosetta2 (DE3) cells and purified from inclusion bodies using fast  
472 protein liquid chromatography as described previously (44, 64).

473

474   **Real-Time Quaking Induced Conversion (RT-QuIC) assay for brain and lymphoid**  
475   **tissue samples**

476           We included brain and lymphoid tissue homogenates from clinical CWD-affected  
477 white-tailed deer, age group matched non-inoculated pigs, and blank (buffer) as controls.  
478 Samples were collected using a strict aseptic technique to minimize the risk of cross-  
479 contamination. All of the samples were run using a blinded study design (N.K., S.M.).

480           Prior to testing, brain and lymphoid tissue samples were homogenized in 1X PBS  
481 at a concentration of 20% w/v tissue, and then further homogenized using repeated  
482 pipetting and sonication in a cup sonicator with two pulses of 30 seconds. The samples  
483 were then further diluted to a concentration of 0.02% in sample dilution buffer (0.025%  
484 SDS in 1X concentration of PBS).

485           The RT-QuIC assay was performed using previously published protocols (41, 65)  
486 with slight modifications as described previously (64). All samples were run in  
487 quadruplicate. The reactions consisted of 5 µg of protein from the brain and lymphoid  
488 tissue homogenates that were used as seed in a 100 µL total reaction volume. A sample  
489 was considered positive if the fluorescence intensity of at least half the replicate wells  
490 crossed the threshold (Ct), which was calculated as the mean fluorescence of the negative

491 control sample plus 10 standard deviations (66-68). For each positive sample we  
492 quantified the seeding activity based on amyloid formation rate (AFR), which is the  
493 reciprocal of the time (h) that it takes for a reaction to reach the threshold (Ct), defined as  
494 the mean baseline fluorescence plus 5 standard deviations (41, 65). The AFR was  
495 calculated using all 4 replicates of each sample. Data analysis was performed using  
496 Biotek's Gen5 software version 2.07.17 and BMG's MARS software version 5.2.R8.

497

## 498 **Acknowledgements**

499 The authors thank Martha Church, Kevin Hassall, Robyn Kokemuller, Joe Lesan,  
500 Leisa Mandell, Dennis Orcutt, and Trudy Tatum for providing critical technical support  
501 to this project. The PoPrP-Tg002 mice were produced by Dr. Glenn Telling through  
502 specific cooperative agreement 58-3625-7-649 with the University of Kentucky. This  
503 study was carried out under the guidelines of the institutional ACUC committee at  
504 NADC. We are grateful to the APHA Biological Archive for providing tissue from a pig  
505 with BSE for use as a positive control for immunohistochemistry and western blot. We  
506 would also like to acknowledge Dr. Byron Caughey for providing the rPrP construct used  
507 in this study. Mention of trade names or commercial products in this article is solely for  
508 the purpose of providing specific information and does not imply recommendation or  
509 endorsement by the US Department of Agriculture. USDA is an equal opportunity  
510 employer.

## References

1. Jahns H, Callanan JJ, Sammin DJ, McElroy MC, Bassett HF. 2006. Survey for transmissible spongiform encephalopathies in Irish pigs fed meat and bone meal. *Vet Rec* 159:137-42.
2. Köfler M, Seuberlich T, Maurer E, Heim D, Doherr M, Zurbriggen A, Botteron C. 2006. [TSE surveillance in small ruminants and pigs: a pilot study]. *Schweiz Arch Tierheilkd* 148:341-2, 344-8.
3. Gajdusek DC, Gibbs CJ, Jr., Alpers M. 1967. Transmission and passage of experimental "kuru" to chimpanzees. *Science* 155:212-4.
4. Hedman C, Bolea R, Marin B, Cobriere F, Filali H, Vazquez F, Pitarch JL, Vargas A, Acin C, Moreno B, Pumarola M, Andreoletti O, Badiola JJ. 2016. Transmission of sheep-bovine spongiform encephalopathy to pigs. *Vet Res* 47:14.
5. Greenlee JJ, Kunkle RA, Smith JD, West Greenlee MH. 2016. Scrapie in swine: a diagnostic challenge. *Food Safety* 4:110-114.
6. Wells GA, Hawkins SA, Austin AR, Ryder SJ, Done SH, Green RB, Dexter I, Dawson M, Kimberlin RH. 2003. Studies of the transmissibility of the agent of bovine spongiform encephalopathy to pigs. *J Gen Virol* 84:1021-31.
7. Dawson M, Wells GA, Parker BN, Scott AC. 1990. Primary parenteral transmission of bovine spongiform encephalopathy to the pig. *Vet Rec* 127:338.
8. Ryder SJ, Hawkins SA, Dawson M, Wells GA. 2000. The neuropathology of experimental bovine spongiform encephalopathy in the pig. *J Comp Pathol* 122:131-43.



- 534 9. Williams ES, Young S. 1980. Chronic wasting disease of captive mule deer: a  
535 spongiform encephalopathy. J Wildl Dis 16:89-98.
- 536 10. Miller MW, Wild MA. 2004. Epidemiology of chronic wasting disease in captive  
537 white-tailed and mule deer. J Wildl Dis 40:320-7.
- 538 11. Spraker TR, Miller MW, Williams ES, Getzy DM, Adrian WJ, Schoonveld GG,  
539 Spowart RA, O'Rourke KI, Miller JM, Merz PA. 1997. Spongiform  
540 encephalopathy in free-ranging mule deer (*Odocoileus hemionus*), white-tailed  
541 deer (*Odocoileus virginianus*) and Rocky Mountain elk (*Cervus elaphus nelsoni*)  
542 in North Central Colorado. J Wildl Dis 33:1-6.
- 543 12. Williams ES, Young S. 1982. Spongiform encephalopathy of Rocky Mountain  
544 elk. J Wildl Dis 18:465-71.
- 545 13. Baeten LA, Powers BE, Jewell JE, Spraker TR, Miller MW. 2007. A natural case  
546 of chronic wasting disease in a free-ranging moose (*Alces alces shirasi*). J Wildl  
547 Dis 43:309-14.
- 548 14. Kreeger TJ, Montgomery DL, Jewell JE, Schultz W, Williams ES. 2006. Oral  
549 transmission of chronic wasting disease in captive Shira's moose. J Wildl Dis  
550 42:640-5.
- 551 15. Benestad SL, Mitchell G, Simmons M, Ytrehus B, Vikoren T. 2016. First case of  
552 chronic wasting disease in Europe in a Norwegian free-ranging reindeer. Vet Res  
553 47:88.
- 554 16. Balachandran A, Harrington NP, Algire J, Soutyrine A, Spraker TR, Jeffrey M,  
555 Gonzalez L, O'Rourke KI. 2010. Experimental oral transmission of chronic

- 556           wasting disease to red deer (*Cervus elaphus elaphus*): early detection and late  
557           stage distribution of protease-resistant prion protein. Can Vet J 51:169-78.
- 558   17.   Hamir AN, Greenlee JJ, Nicholson EM, Kunkle RA, Richt JA, Miller JM, Hall M.  
559           2011. Experimental transmission of chronic wasting disease (CWD) from elk and  
560           white-tailed deer to fallow deer by intracerebral route: final report. Can J Vet Res  
561           75:152-6.
- 562   18.   Nalls AV, McNulty E, Powers J, Seelig DM, Hoover C, Haley NJ, Hayes-Klug J,  
563           Anderson K, Stewart P, Goldmann W, Hoover EA, Mathiason CK. 2013. Mother  
564           to offspring transmission of chronic wasting disease in reeves' muntjac deer.  
565           PLoS One 8:e71844.
- 566   19.   Moore SJ, Kunkle R, Greenlee MH, Nicholson E, Richt J, Hamir A, Waters WR,  
567           Greenlee J. 2016. Horizontal transmission of chronic wasting disease in reindeer.  
568           Emerg Infect Dis 22:2142-2145.
- 569   20.   Mitchell GB, Sigurdson CJ, O'Rourke KI, Algire J, Harrington NP, Walther I,  
570           Spraker TR, Balachandran A. 2012. Experimental oral transmission of chronic  
571           wasting disease to reindeer (*Rangifer tarandus tarandus*). PLoS One 7:e39055.
- 572   21.   Hamir AN, Kunkle RA, Cutlip RC, Miller JM, Williams ES, Richt JA. 2006.  
573           Transmission of chronic wasting disease of mule deer to Suffolk sheep following  
574           intracerebral inoculation. J Vet Diagn Invest 18:558-65.
- 575   22.   Greenlee JJ, Nicholson EM, Smith JD, Kunkle RA, Hamir AN. 2012.  
576           Susceptibility of cattle to the agent of chronic wasting disease from elk after  
577           intracranial inoculation. J Vet Diagn Invest 24:1087-93.

- 578 23. Hamir AN, Cutlip RC, Miller JM, Williams ES, Stack MJ, Miller MW, O'Rourke  
579 KI, Chaplin MJ. 2001. Preliminary findings on the experimental transmission of  
580 chronic wasting disease agent of mule deer to cattle. *J Vet Diagn Invest* 13:91-6.
- 581 24. Hamir AN, Kunkle RA, Cutlip RC, Miller JM, O'Rourke KI, Williams ES, Miller  
582 MW, Stack MJ, Chaplin MJ, Richt JA. 2005. Experimental transmission of  
583 chronic wasting disease agent from mule deer to cattle by the intracerebral route. *J*  
584 *Vet Diagn Invest* 17:276-81.
- 585 25. Hamir AN, Kunkle RA, Miller JM, Greenlee JJ, Richt JA. 2006. Experimental  
586 second passage of chronic wasting disease (CWD(mule deer)) agent to cattle. *J*  
587 *Comp Pathol* 134:63-9.
- 588 26. Mathiason CK, Nalls AV, Seelig DM, Kraft SL, Carnes K, Anderson KR, Hayes-  
589 Klug J, Hoover EA. 2013. Susceptibility of domestic cats to chronic wasting  
590 disease. *J Virol* 87:1947-56.
- 591 27. Seelig DM, Nalls AV, Flasik M, Frank V, Eaton S, Mathiason CK, Hoover EA.  
592 2015. Lesion profiling and subcellular prion localization of cervid chronic  
593 wasting disease in domestic cats. *Vet Pathol* 52:107-19.
- 594 28. Bartz JC, Marsh RF, McKenzie DI, Aiken JM. 1998. The host range of chronic  
595 wasting disease is altered on passage in ferrets. *Virology* 251:297-301.
- 596 29. Sigurdson CJ, Mathiason CK, Perrott MR, Eliason GA, Spraker TR, Glatzel M,  
597 Manco G, Bartz JC, Miller MW, Hoover EA. 2008. Experimental chronic wasting  
598 disease (CWD) in the ferret. *J Comp Pathol* 138:189-96.

- 599 30. Marsh RF, Kincaid AE, Bessen RA, Bartz JC. 2005. Interspecies transmission of  
600 chronic wasting disease prions to squirrel monkeys (*Saimiri sciureus*). J Virol  
601 79:13794-6.
- 602 31. Race B, Meade-White KD, Miller MW, Barbian KD, Rubenstein R, LaFauci G,  
603 Cervenakova L, Favara C, Gardner D, Long D, Parnell M, Striebel J, Priola SA,  
604 Ward A, Williams ES, Race R, Chesebro B. 2009. Susceptibilities of nonhuman  
605 primates to chronic wasting disease. Emerg Infect Dis 15:1366-76.
- 606 32. Race B, Meade-White KD, Phillips K, Striebel J, Race R, Chesebro B. 2014.  
607 Chronic wasting disease agents in nonhuman primates. Emerg Infect Dis 20:833-  
608 7.
- 609 33. Kurt TD, Sigurdson CJ. 2016. Cross-species transmission of CWD prions. Prion  
610 10:83-91.
- 611 34. US Department of Agriculture National Feral Swine Mapping System. 2015.  
612 <http://swine.vet.uga.edu/nfsms/index.jsp>. Accessed Feb 13.
- 613 35. US Geological Survey National Wildlife Health Center. 2017.  
614 [https://www.nwhc.usgs.gov/disease\\_information/chronic\\_wasting\\_disease/index.j](https://www.nwhc.usgs.gov/disease_information/chronic_wasting_disease/index.jsp)  
615 [sp](https://www.nwhc.usgs.gov/disease_information/chronic_wasting_disease/index.jsp). Accessed Feb 13.
- 616 36. Pritzkow S, Morales R, Moda F, Khan U, Telling GC, Hoover E, Soto C. 2015.  
617 Grass plants bind, retain, uptake, and transport infectious prions. Cell Rep  
618 11:1168-75.
- 619 37. Johnson CJ, Phillips KE, Schramm PT, McKenzie D, Aiken JM, Pedersen JA.  
620 2006. Prions adhere to soil minerals and remain infectious. PLoS Pathog 2:e32.

- 621 38. Johnson CJ, Pedersen JA, Chappell RJ, McKenzie D, Aiken JM. 2007. Oral  
622 transmissibility of prion disease is enhanced by binding to soil particles. *PLoS*  
623 *Pathog* 3:e93.
- 624 39. Seidel B, Thomzig A, Buschmann A, Groschup MH, Peters R, Beekes M, Terytze  
625 K. 2007. Scrapie Agent (Strain 263K) can transmit disease via the oral route after  
626 persistence in soil over years. *PLoS One* 2:e435.
- 627 40. Rodriguez-Martinez AB, Garrido JM, Zarranz JJ, Arteagoitia JM, de Pancorbo  
628 MM, Atares B, Bilbao MJ, Ferrer I, Juste RA. 2010. A novel form of human  
629 disease with a protease-sensitive prion protein and heterozygosity  
630 methionine/valine at codon 129: Case report. *BMC Neurol* 10:99.
- 631 41. Wilham JM, Orru CD, Bessen RA, Atarashi R, Sano K, Race B, Meade-White  
632 KD, Taubner LM, Timmes A, Caughey B. 2010. Rapid end-point quantitation of  
633 prion seeding activity with sensitivity comparable to bioassays. *PLoS Pathog*  
634 6:e1001217.
- 635 42. Dassanayake RP, Orru CD, Hughson AG, Caughey B, Graca T, Zhuang D,  
636 Madsen-Bouterse SA, Knowles DP, Schneider DA. 2016. Sensitive and specific  
637 detection of classical scrapie prions in the brains of goats by real-time quaking-  
638 induced conversion. *J Gen Virol* 97:803-12.
- 639 43. Peden AH, McGuire LI, Appleford NE, Mallinson G, Wilham JM, Orru CD,  
640 Caughey B, Ironside JW, Knight RS, Will RG, Green AJ, Head MW. 2012.  
641 Sensitive and specific detection of sporadic Creutzfeldt-Jakob disease brain prion  
642 protein using real-time quaking-induced conversion. *J Gen Virol* 93:438-49.

- 643 44. Kondru N, Manne S, Greenlee J, West Greenlee H, Anantharam V, Halbur P,  
644 Kanthasamy A, Kanthasamy A. 2017. Integrated organotypic slice cultures and  
645 RT-QuIC (OSCAR) assay: implications for translational discovery in protein  
646 misfolding diseases. *Sci Rep* 7:43155.
- 647 45. Vascellari S, Orru CD, Hughson AG, King D, Barron R, Wilham JM, Baron GS,  
648 Race B, Pani A, Caughey B. 2012. Prion seeding activities of mouse scrapie  
649 strains with divergent PrP<sup>Sc</sup> protease sensitivities and amyloid plaque content  
650 using RT-QuIC and eQuIC. *PLoS One* 7:e48969.
- 651 46. Tzaban S, Friedlander G, Schonberger O, Horonchik L, Yedidia Y, Shaked G,  
652 Gabizon R, Taraboulos A. 2002. Protease-sensitive scrapie prion protein in  
653 aggregates of heterogeneous sizes. *Biochemistry* 41:12868-75.
- 654 47. Pastrana MA, Sajnani G, Onisko B, Castilla J, Morales R, Soto C, Requena JR.  
655 2006. Isolation and characterization of a proteinase K-sensitive PrP(Sc) fraction.  
656 *Biochemistry* 45:15710-15717.
- 657 48. Mathiason CK, Powers JG, Dahmes SJ, Osborn DA, Miller KV, Warren RJ,  
658 Mason GL, Hays SA, Hayes-Klug J, Seelig DM, Wild MA, Wolfe LL, Spraker  
659 TR, Miller MW, Sigurdson CJ, Telling GC, Hoover EA. 2006. Infectious prions  
660 in the saliva and blood of deer with chronic wasting disease. *Science* 314:133-6.
- 661 49. Mathiason CK, Hays SA, Powers J, Hayes-Klug J, Langenberg J, Dahmes SJ,  
662 Osborn DA, Miller KV, Warren RJ, Mason GL, Hoover EA. 2009. Infectious  
663 prions in pre-clinical deer and transmission of chronic wasting disease solely by  
664 environmental exposure. *PLoS One* 4:e5916.

- 665 50. Haley NJ, Seelig DM, Zabel MD, Telling GC, Hoover EA. 2009. Detection of  
666 CWD prions in urine and saliva of deer by transgenic mouse bioassay. PLoS ONE  
667 4:e4848.
- 668 51. Henderson DM, Manca M, Haley NJ, Denkers ND, Nalls AV, Mathiason CK,  
669 Caughey B, Hoover EA. 2013. Rapid antemortem detection of CWD prions in  
670 deer saliva. PLoS One 8:e74377.
- 671 52. Haley NJ, Mathiason CK, Zabel MD, Telling GC, Hoover EA. 2009. Detection of  
672 sub-clinical CWD infection in conventional test-negative deer long after oral  
673 exposure to urine and feces from CWD+ deer. PLoS One 4:e7990.
- 674 53. Tamguney G, Miller MW, Wolfe LL, Sirochman TM, Glidden DV, Palmer C,  
675 Lemus A, DeArmond SJ, Prusiner SB. 2009. Asymptomatic deer excrete  
676 infectious prions in faeces. Nature 461:529-32.
- 677 54. Pulford B, Spraker TR, Wyckoff AC, Meyerett C, Bender H, Ferguson A, Wyatt  
678 B, Lockwood K, Powers J, Telling GC, Wild MA, Zabel MD. 2012. Detection of  
679 PrPCWD in feces from naturally exposed Rocky Mountain elk (*Cervus elaphus*  
680 *nelsoni*) using protein misfolding cyclic amplification. J Wildl Dis 48:425-34.
- 681 55. Wilcox J, Van Vuren D. 2009. Wild pigs as predators in oak woodlands of  
682 California. J Mammal 90:114-118.
- 683 56. Hamir AN, Richt JA, Miller JM, Kunkle RA, Hall SM, Nicholson EM, O'Rourke  
684 KI, Greenlee JJ, Williams ES. 2008. Experimental transmission of chronic  
685 wasting disease (CWD) of elk (*Cervus elaphus nelsoni*), white-tailed deer  
686 (*Odocoileus virginianus*), and mule deer (*Odocoileus hemionus hemionus*) to  
687 white-tailed deer by intracerebral route. Vet Pathol 45:297-306.

- 688 57. Greenlee JJ, Smith JD, Kunkle RA. 2011. White-tailed deer are susceptible to the  
689 agent of sheep scrapie by intracerebral inoculation. *Vet Res* 42:107.
- 690 58. Smith JD, Nicholson EM, Greenlee JJ. 2013. Evaluation of a combinatorial  
691 approach to prion inactivation using an oxidizing agent, SDS, and proteinase K.  
692 *BMC Vet Res* 9:151.
- 693 59. Smith JD, Nicholson EM, Foster GH, Greenlee JJ. 2013. Exposure of RML  
694 scrapie agent to a sodium percarbonate-based product and sodium dodecyl sulfate  
695 renders PrPSc protease sensitive but does not eliminate infectivity. *BMC Vet Res*  
696 9:8.
- 697 60. Greenlee JJ, Smith JD, West Greenlee MH, Nicholson EM. 2012. Clinical and  
698 pathologic features of H-type bovine spongiform encephalopathy associated with  
699 E211K prion protein polymorphism. *PLoS ONE* 7:e38678.
- 700 61. Smith JD, Greenlee JJ. 2014. Detection of misfolded prion protein in retina  
701 samples of sheep and cattle by use of a commercially available enzyme  
702 immunoassay. *Am J Vet Res* 75:268-72.
- 703 62. Hardt M, Baron T, Groschup MH. 2000. A comparative study of  
704 immunohistochemical methods for detecting abnormal prion protein with  
705 monoclonal and polyclonal antibodies. *J Comp Pathol* 122:43-53.
- 706 63. Atarashi R, Wilham JM, Christensen L, Hughson AG, Moore RA, Johnson LM,  
707 Onwubiko HA, Priola SA, Caughey B. 2008. Simplified ultrasensitive prion  
708 detection by recombinant PrP conversion with shaking. *Nat Methods* 5:211-2.
- 709 64. West Greenlee MH, Lind M, Kokemueller R, Mammadova N, Kondru N, Manne  
710 S, Smith J, Kanthasamy A, Greenlee J. 2016. Temporal resolution of misfolded



- 711 prion protein transport, accumulation, glial activation, and neuronal death in the  
712 retinas of mice inoculated with scrapie. *Am J Pathol* 186:2302-9.
- 713 65. Orru CD, Groveman BR, Hughson AG, Zanusso G, Coulthart MB, Caughey B.  
714 2015. Rapid and sensitive RT-QuIC detection of human Creutzfeldt-Jakob disease  
715 using cerebrospinal fluid. *MBio* 6.
- 716 66. Groveman BR, Orru CD, Hughson AG, Bongianni M, Fiorini M, Imperiale D,  
717 Ladogana A, Pocchiari M, Zanusso G, Caughey B. 2017. Extended and direct  
718 evaluation of RT-QuIC assays for Creutzfeldt-Jakob disease diagnosis. *Ann Clin*  
719 *Transl Neurol* 4:139-144.
- 720 67. Henderson DM, Davenport KA, Haley NJ, Denkers ND, Mathiason CK, Hoover  
721 EA. 2015. Quantitative assessment of prion infectivity in tissues and body fluids  
722 by real-time quaking-induced conversion. *J Gen Virol* 96:210-9.
- 723 68. Bongianni M, Orru C, Groveman BR, Sacchetto L, Fiorini M, Tonoli G, Triva G,  
724 Capaldi S, Testi S, Ferrari S, Cagnin A, Ladogana A, Poleggi A, Colaizzo E,  
725 Tiple D, Vaianella L, Castriciano S, Marchioni D, Hughson AG, Imperiale D,  
726 Cattaruzza T, Fabrizi GM, Pocchiari M, Monaco S, Caughey B, Zanusso G. 2017.  
727 Diagnosis of human prion disease using real-time quaking-induced conversion  
728 testing of olfactory mucosa and cerebrospinal fluid samples. *JAMA Neurol*  
729 74:155-162.
- 730

731 **Tables**732 **Table 1. Detection and characterization of disease-associated prion protein (PrP<sup>Sc</sup>) from selected pigs.**

Animal number	Treatment group	Incubation period <sup>a</sup>	Overall result CNS	Overall result LRS	Antigen-capture enzyme immunoassay	Western blot	Immuno-histochemistry	Proteinase-K sensitivity	RT-QuIC result	RT-QuIC amyloid formation rate
1	Control market weight	0	-	nt	na	na	nt	nt	nt	nt
2		0	-	nt	-	-	nt	nt	nt	nt
3		0	-	nt	-	-	nt	nt	nt	nt
4		6	-	nt	-	-	nt	nt	nt	nt
5		6	-	-	-	-	-	na	-	0
6	Control aged	25	-	nt	-	-	nt	nt	nt	nt
7		41	-	nt	-	-	nt	nt	nt	nt
8		46	-	nt	-	-	nt	nt	nt	nt
9		73	-	-	-	-	-	na	-	0
10	Intracranially inoculated market weight	0	-	nt	-	-	nt	nt	nt	nt
11		0	-	nt	-	-	nt	nt	nt	nt
12		6	+	+	-	-	-	na	+	0.031
13		6	-	nt	-	-	nt	nt	nt	nt
14		6	+	+	-	-	-	na	+	0.025
15		6	+	+	+	-	-	sensitive	-	0
16		6	-	+	-	-	-	na	-	0
17		6	-	nt	-	-	nt	nt	nt	nt
18		6	+	+	-	-	-	na	+	0.120
19		6	-	-	-	-	-	na	-	0
20	Intracranially inoculated aged	30	-	nt	-	-	nt	nt	nt	nt
21		30	-	nt	-	-	nt	nt	nt	nt
22		30	+	+	-	-	-	na	+	0.080

23		30	-	nt	-	-	nt	nt	nt	nt
24		42	+	+	+	-	-	sensitive	+	0.030
25		45	+	+	+	-	+	resistant	+	0.180
26		56	+	+	+	-	+	resistant	+	0.190
27		64	+	-	+	+	-	resistant	+	0.170
28		73	+	+	+	+	+	resistant	+	0.210
29		73	+	+	-	-	-	na	+	0.050
32		6	+	+	-	-	-	na	+	0.070
38		6	+	+	-	-	-	na	+	0.010
30		6	-	+	-	-	-	na	-	0
36		6	-	+	-	-	-	na	-	0
37		6	-	+	-	-	-	na	-	0
34		6	-	-	-	-	-	na	-	0
31		6	-	nt	-	-	nt	nt	nt	nt
33		6	-	nt	-	-	nt	nt	nt	nt
35		6	-	nt	-	-	nt	nt	nt	nt
39		19	-	+	-	-	-	na	-	0
40		41	-	nt	-	-	nt	nt	nt	nt
41		41	+	+	-	-	-	na	+	0.029
42		41	-	nt	-	-	nt	nt	nt	nt
43		45	+	+	-	-	-	na	+	0.020
44		55	+	+	-	-	-	na	+	0.030
45		64	+	-	+	-	+	sensitive	+	0.010
46		65	-	nt	-	-	nt	nt	nt	nt
47		65	-	nt	-	-	nt	nt	nt	nt
48		72	+	-	-	-	-	na	+	0.010

733 Incubation periods are expressed as months post-inoculation. RT-QuIC, real-time quaking-induced conversion assay; na, result not  
734 applicable; nt, sample not tested.

735 **Table 2. Detection of disease-associated prion protein (PrP<sup>Sc</sup>) in lymphoid tissues using antigen-capture enzyme immunoassay**  
 736 **(EIA) and real-time quaking induced conversion assay (RT-QuIC).**

Animal number	Treatment group	Incubation period <sup>a</sup>	Overall result	Retropharyngeal lymph node		Tonsil		Mesenteric lymph node	
				EIA	RT-QuIC	EIA	RT-QuIC	EIA	RT-QuIC
5	Non-inoculated controls	6	-	-	-	-	-	-	-
9		73	-	na	na	-	-	-	-
12	Intracranially inoculated market weight	6	+	-	-	-	-	-	+
14		6	+	-	+	-	+	-	-
15		6	+	-	+	-	+	-	+
16		6	+	-	+	-	+	-	+
18		6	+	-	-	-	-	-	+
19		6	-	-	-	-	-	-	-
22	Intracranially inoculated aged	30	+	-	-	-	-	-	+
24		42	+	na	na	-	+	na	na
25		45	+	-	+	-	-	-	-
26		56	+	-	+	-	+	na	na
27		64	-	na	na	-	-	na	na
28		73	+	na	na	-	+	-	+
29	Orally inoculated market weight	73	+	-	+	-	-	-	-
30		6	+	-	-	-	-	-	+
32		6	+	-	-	-	+	-	+
34		6	-	-	-	-	-	-	-
36		6	+	-	-	-	+	-	+
37		6	+	-	+	-	-	+	+
38	Orally inoculated aged	6	+	-	-	-	+	-	+
39		19	+	-	-	-	-	-	+
41		41	+	-	+	-	-	-	+
43		45	+	-	na	-	+	na	na
44		55	+	na	na	-	-	-	+
45		64	-	-	na	-	-	na	na
48		72	-	na	na	-	-	-	-

737 <sup>a</sup> Incubation periods are expressed as months post-inoculation. na, sample not available.

738 **Table 3. Results from bioassay of brain material from selected pigs in *Tg002* mice**  
 739 **that express porcine prion protein.**

740

Donor animal number	Donor treatment group (donor incubation period <sup>a</sup> )	<i>Tg002</i> attack rate <sup>b</sup>	<i>Tg002</i> mean incubation period <sup>c</sup>
18	Intracranially inoculated market weight (6)	2/29	244
27	Intracranially inoculated aged (64)	3/29	167
28	Intracranially inoculated aged (73)	2/27	314, 701 <sup>d</sup>
32	Orally inoculated market weight (6)	0/28	>700
48	Orally inoculated aged (72)	14/28	263

741

742 <sup>a</sup> Donor incubation period is expressed as months post-inoculation. <sup>b</sup> Disease-associated  
 743 prion protein in the brains of mice was detected using an antigen-capture enzyme  
 744 immunoassay (EIA). <sup>c</sup> Mouse incubation period is expressed as days post-inoculation.  
 745 Survival <sup>d</sup> Survival times for these 2 mice are so disparate that calculation of a mean  
 746 incubation period would not be meaningful.

747

748 **Figure legends**

749

750 **Figure 1. Western blot analysis demonstrating a unique PrP<sup>Sc</sup> profile in brain**  
751 **samples from pigs with CWD.**

752 The positive brain sample from a pig inoculated with the CWD agent (Pig CWD) has a  
753 slightly higher migration relative to a brain sample from a pig inoculated with the agent  
754 of classical bovine spongiform encephalopathy (Pig BSE), and a much lower migration  
755 relative to the CWD inoculum (CWD Inoc). The diglycosylated band (top most band in  
756 each lane) is more prominent in the Pig CWD and CWD Inoc samples, while the  
757 monoglycosylated (middle) band is most prominent in the Pig BSE sample. Blot  
758 developed with monoclonal antibody L42. Note: due to the sparse PrP<sup>Sc</sup> accumulation in  
759 the brains of inoculated pigs the blot in Figure 1 is a composite, see Materials and  
760 methods for details.

761

762

763 **Figure 2. Amyloid formation rates (RT-QuIC) and proteinase-K sensitivity (EIA) of**  
764 **PrP<sup>Sc</sup> from pig brain samples.**

765 Treatment groups: animals 5 and 9, non-inoculated controls; 12-19, IC inoculated market  
766 weight pigs; 22-29, IC inoculated aged pigs; 30-38, orally inoculated market weight pigs;  
767 39-48, orally inoculated aged pigs. Proteinase-K (PK) sensitivity: solid fill, PK sensitivity  
768 not determined (EIA negative); horizontal stripe fill, PK resistant; checked fill, PK  
769 sensitive.

770

771

772 **Figure 3. Results of RT-QuIC assays of brain homogenate from inoculated and**  
773 **negative control pigs.**

774 Average percent thioflavin T (ThT) fluorescence readings (thick lines) with standard  
775 deviations (thin lines) determined from all replicates (four replicate reactions per animal)  
776 from all pigs in each challenge group. Red, intracranially inoculated aged pigs (n = 7);  
777 blue, intracranially inoculated market-weight pigs (n = 6); purple, orally inoculated  
778 market-weight pigs (n = 6); orange, orally inoculated aged pigs (n = 6); green, non-  
779 inoculated control pigs (n = 2).

782 **Figure 4. Vacuolar change and disease-associated prion protein in the brain and**  
783 **eye.**

784 (A) Brainstem (Pig #7). Incidental, i.e. not related to prion disease, neuropil vacuolation in  
785 the colliculus, \* midline (hematoxylin and eosin, original magnification 4x). (B) Higher  
786 magnification view of (A) (original magnification 10x). (C) Brainstem (Pig #25).  
787 Intraneuronal PrP<sup>Sc</sup> immunoreactivity (arrows) in neurons in the colliculus (monoclonal  
788 anti-PrP antibody L42, original magnification 20x). (D) Brainstem (Pig #8). Non disease-  
789 specific intraneuronal immunolabeling (arrows) in neurons in the colliculus from a non-  
790 inoculated control pig (monoclonal anti-PrP antibody L42, original magnification 40x).  
791 (E) Brainstem (Pig #38). Incidental intraneuronal vacuolation (\*) in the dorsal motor  
792 nucleus of the vagus nerve (hematoxylin and eosin, original magnification 40x). (F)  
793 Retina (Pig #26). Granular to punctate PrP<sup>Sc</sup> immunoreactivity in the inner and out  
794 plexiform layers with occasional intragial deposits (arrow) (monoclonal anti-PrP

795 antibody L42, original magnification 40x). (G) Retina (Pig #4). Non disease-specific  
796 immunolabeling in a non-inoculated control pig (monoclonal anti-PrP antibody L42,  
797 original magnification 40x).









